

A Uniqueness Property for H^∞ on Coverings of Projective Manifolds

Alexander Brudnyi*

Department of Mathematics and Statistics
University of Calgary
Calgary, Canada

Abstract

Let Y be a regular covering of a complex projective manifold $M \hookrightarrow \mathbb{CP}^N$ of dimension $n \geq 2$. Let C be intersection with M of at most $n - 1$ generic hypersurfaces of degree d in $\mathbb{CP}^N$. The preimage X of C in Y is a connected submanifold. Let $H^\infty(Y)$ and $H^\infty(X)$ be the Banach spaces of bounded holomorphic functions on Y and X in the corresponding supremum norms. We prove that the restriction $H^\infty(Y) \longrightarrow H^\infty(X)$ is an isometry for d large enough. This answers the question posed in [L].

1. Introduction and Formulation of the Result.

1.1. An Extension Theorem. Let M be a complex projective manifold of dimension $n \geq 2$ with a Kähler form ω and let L be a positive line bundle on M with canonical connection ∇ and curvature Θ in a hermitian metric h . Let C be the common zero locus of holomorphic sections $s_1, \dots, s_k$, $k < n$, of L over M which, in a trivialization, can be completed to a set of local coordinates at each point C . Then C is a (possibly disconnected) k -dimensional submanifold of M which will be referred to as an *L -submanifold of M* . Let $\pi : Y_G \longrightarrow M$ be a regular covering of M with a transformation group G and $X_G = \pi^{-1}(C)$. We denote the pullbacks to Y_G of ω and Θ by the same letters.

Example 1.1 If L is very ample, then it is pullback of the hyperplane bundle by an embedding of M into some projective space $\mathbb{CP}^N$. Further, zero loci of holomorphic sections of L are hyperplane sections of M . By Bertini's theorem, the generic linear subspace of codimension $n-k$, $k < n$, intersects M transversely in a smooth manifold C of dimension k , and by the Lefschetz hyperplane theorem, C is connected and

*Research supported in part by NSERC.

2000 *Mathematics Subject Classification*. Primary 32A10. Secondary 14E20.

Key words and phrases. Bounded holomorphic function, corona problem, transformation group.

the induced homomorphism $\pi_1(C) \longrightarrow \pi_1(M)$ of fundamental groups is surjective. Thus in this case $X_G \subset Y_G$ is a connected submanifold.

Further, let $\text{dist}(\cdot, \cdot)$ be the distance on Y_G induced by ω . Consider a function $\phi : Y_G \longrightarrow \mathbb{R}$ such that $d\phi$ is bounded, i.e.

$$|\phi(x) - \phi(y)| \leq a \cdot \text{dist}(x, y) \quad \text{for some } a > 0.$$

By $\mathcal{O}_\phi(X_G)$ we denote the vector space of holomorphic functions on X_G such that $|f|^2 e^{-\phi}$ is integrable on X_G with respect to the volume form of the induced Kähler metric on X_G . This is a Hilbert space with respect to the inner product

$$(f, g) \mapsto \int_{X_G} f \bar{g} e^{-\phi} \omega^k.$$

We define $\mathcal{O}_\phi(Y_G)$ similarly. By $|\cdot|_{\phi, X_G}$ and $|\cdot|_{\phi, Y_G}$ we denote the corresponding norms. It was shown in [L] that the restriction determines a continuous linear map

$$\rho : \mathcal{O}_\phi(Y_G) \longrightarrow \mathcal{O}_\phi(X_G), \quad f \mapsto f|_{X_G}.$$

The following remarkable result was proved by Lárusson [L, Th.1.2].

Theorem 1.2 *Suppose*

$$\Theta \geq i\partial\bar{\partial}\phi + \epsilon\omega$$

for some $\epsilon > 0$ in the sense of Nakano. Then ρ is an isomorphism.

1.2. Formulation of the Main Result. An important example of a function ϕ as above is obtained by smoothing the distance δ from a fixed point o in Y_G . By a result of Napier [N], there is a smooth function τ on Y_G such that

$$(A) \quad c_1\delta \leq \tau \leq c_2\delta + c_3 \text{ for some } c_1, c_2, c_3 > 0,$$

$$(B) \quad d\tau \text{ is bounded, and}$$

$$(C) \quad i\partial\bar{\partial}\tau \text{ is bounded.}$$

Furthermore, by (A) and since the curvature of Y_G is bounded below, there is $c > 0$ such that $e^{-c\tau}$ is integrable on Y_G . Then $e^{-c\tau}$ is also integrable on X_G . We set

$$A := \frac{cc_2}{c_1}. \tag{1.1}$$

Let $\tilde{L}$ be any positive line bundle on M with curvature $\tilde{\Theta}$. By (C) there is a non-negative integer m_0 such that for any integer $m > m_0$

$$m\tilde{\Theta} > i\partial\bar{\partial}(A\tau). \tag{1.2}$$

We set $L := \tilde{L}^{\otimes m}$. Then Lárusson's theorem holds for coverings $X_G := \pi^{-1}(C) \subset Y_G$ of L -submanifolds $C \subset M$ with $\phi := A\tau$ and with $\phi := c\tau$ (because $A \geq c$).

Let $H^\infty(Y_G)$, $H^\infty(X_G)$ be the Banach spaces of bounded holomorphic functions on Y_G and X_G in the corresponding supremum norms.

Theorem 1.3 *The map $\rho : H^\infty(Y_G) \longrightarrow H^\infty(X_G)$, $f \mapsto f|_{X_G}$, is an isometry.*

1.3. Corollaries and Examples. Let X be a complex manifold and $H^\infty(X)$ be the Banach algebra (in the supremum norm) of bounded holomorphic functions on X . The maximal ideal space $\mathcal{M} = \mathcal{M}(H^\infty(X))$ is the set of all nontrivial linear multiplicative functionals on $H^\infty(X)$. The norm of any $\phi \in \mathcal{M}$ is ≤ 1 and so $\mathcal{M}$ is embedded into the unit ball of the dual space $(H^\infty(X))^*$. Then $\mathcal{M}$ is a compact Hausdorff space in the weak $*$ topology induced by $(H^\infty(X))^*$ (i.e. the *Gelfand topology*). Further, there is a continuous map $i : X \longrightarrow \mathcal{M}$ taking $x \in X$ to the evaluation homomorphism $f \mapsto f(x)$. This map is an embedding if $H^\infty(X)$ separates points of X . Recall also that the complement to the closure of $i(X)$ in $\mathcal{M}$ is called *the corona*. The *corona problem* is to determine those X for which the corona is empty. For example, according to Carleson's celebrated Corona Theorem [C] this is true for X being the open unit disk $\mathbb{D} \subset \mathbb{C}$. Also there are non-planar Riemann surfaces for which the corona is non-trivial (see e.g. [JM], [G], [BD] and references therein). The general problem for planar domains is still open, as is the problem in several variables for the ball and polydisk. In [L, Th. 2.1] Lárusson discovered simplest examples of Riemann surfaces with big corona. Namely, using his Theorem 1.2 he proved that if $Y_G \subset \mathbb{C}^n$ is a bounded domain and $X_G \subset Y_G$ is a Riemann surface satisfying assumptions of Theorem 1.3 then the natural map $i : X_G \hookrightarrow \mathcal{M}(H^\infty(X_G))$ extends to an embedding $Y_G \hookrightarrow \mathcal{M}(H^\infty(X_G))$. The next statement is an extension of his result.

Corollary 1.4 *Under the assumptions of Theorem 1.3, the transpose map $\rho^* : \mathcal{M}(H^\infty(X_G)) \longrightarrow \mathcal{M}(H^\infty(Y_G))$, $\phi \mapsto \phi \circ \rho$, is a homeomorphism.*

This follows from the fact that $\rho : H^\infty(Y_G) \longrightarrow H^\infty(X_G)$ is an isometry of Banach algebras. $\square$

Example 1.5 (1) (The references for this example are in [L, Sect. 4].) Let M be a projective manifold covered by the unit ball $\mathbb{B} \subset \mathbb{C}^n$ with a positive line bundle L with curvature Θ , and $X \subset \mathbb{B}$ be the preimage of an L -submanifold $C \subset M$. Let δ be the distance from the origin in the Bergman metric of $\mathbb{B}$. By ω we denote the Kähler form of the Bergman metric. It was shown in [L, Sect. 4] that there is a nonnegative function τ on $\mathbb{B}$ such that $i\partial\bar{\partial}\tau = \omega$, $d\tau$ is bounded, and

$$\sqrt{n+1}\delta \leq \tau \leq \sqrt{n+1}\delta + (n+1)\log 2.$$

Moreover,

$$\int_{\mathbb{B}} e^{-c\tau} \omega^n < \infty \quad \text{if and only if} \quad c > \frac{2n}{n+1}.$$

Applying Theorem 1.3 (with $c_2 = c_1 = \sqrt{n+1}$) we obtain that $\rho : H^\infty(\mathbb{B}) \rightarrow H^\infty(X)$ is an isometry if $\Theta > \frac{2n}{n+1}\omega$. This holds for instance if $L = K^{\otimes m}$ with $m \geq 2$ where K is the canonical bundle of M .

(2) Let S be a compact complex curve of genus $g \geq 2$ and $\mathbb{CT}$ be a one-dimensional complex torus. Consider an L -curve C in $M := S \times \mathbb{CT}$ with a very ample L satisfying the assumptions of Theorem 1.3. Let $\pi : \mathbb{D} \times \mathbb{C} \longrightarrow M$ be the universal

covering. Then Theorem 1.3 is valid for the connected curve $X := \pi^{-1}(C) \subset \mathbb{D} \times \mathbb{C}$. This implies that any $f \in H^\infty(X)$ is constant on each $S_y := (\{y\} \times \mathbb{C}) \cap X$, $y \in \mathbb{D}$. Note that S_y is union of the orbits of some $z_{iy} \in X$, $i = 1, \dots, k$, under the natural action of the group $\pi_1(\mathbb{CT})$ ($\cong \mathbb{Z} \oplus \mathbb{Z}$) on $\mathbb{D} \times \mathbb{C}$.

2. Proof of Theorem 1.3.

Let us fix a fundamental compact K of the action of G on Y_G , i.e., $Y_G = \cup_{g \in G} g(K)$. Consider finite covers $\mathcal{U} = (U_i)$ and $\mathcal{V} = (V_j)$ of K by compact co-ordinate polydisks such that each V_j belongs to the interior of some U_{i_j} .

Lemma 2.1 *Let f be a holomorphic function defined in an open neighbourhood O of $\cup_i U_i$. Assume that*

$$\int_O |f|^2 \omega^n = B < \infty .$$

Then there is a constant $b > 0$ (depending on $\mathcal{U}$ and $\mathcal{V}$ only) such that

$$\max_K |f| \leq b\sqrt{B} . \quad (2.1)$$

The proof of the lemma is the consequence of the following facts:

- (a) after the identification of U_{i_j} with the closed unit polydisk D and of V_j with a compact subset $D_j \subset D$, the volume form ω^n restricted to each U_{i_j} is equivalent to the Euclidean volume form $do := dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_n$;
- (b) the Bergman inequality (see [GR, Ch.6, Th.1.3])

$$\max_{D_j} |f| \leq \frac{(\sqrt{n})^n}{(\sqrt{\pi}d)^n} \cdot \left(\int_D |f|^2 do \right)^{1/2} ,$$

where d is the Euclidean distance from D_j to the boundary of D .

We leave the details to the reader. $\square$

Let $\text{dist}(\cdot, \cdot)$ be the distance on Y_G in the metric induced by ω . In particular, $\delta(x) := \text{dist}(x, o)$. Since ω is invariant with respect to the action of G we also have $\text{dist}(g(x), g(y)) = \text{dist}(x, y)$ for any $g \in G$. From inequalities (A) for τ and the triangle inequality for the distance we obtain

$$\begin{aligned} \tau(g(x)) &\geq c_1 \text{dist}(g(x), o) \geq c_1 [\text{dist}(g(x), g(o)) - \text{dist}(g(o), o)] = \\ &c_1 [\text{dist}(x, o) - \text{dist}(g(o), o)] \geq (c_1/c_2)\tau(x) - (c_1 c_3/c_2) - c_1 \delta(g(o)) . \end{aligned} \quad (2.2)$$

Further, if $x \in K$ then

$$a_1 \leq \tau(x) \leq a_2 \quad \text{for some } a_1, a_2 > 0 . \quad (2.3)$$

Below by $|\cdot|_{\infty, X_G}$ and $|\cdot|_{\infty, Y_G}$ we denote the corresponding H^∞ -norms. Let $f \in H^\infty(X_G)$. Then $f \in \mathcal{O}_{A\tau}(X_G) \cap \mathcal{O}_{c\tau}(X_G)$ and there is $a_3 > 0$ such that

$$\max\{|f|_{A\tau, X_G}, |f|_{c\tau, X_G}\} \leq a_3 \sup_{X_G} |f| := a_3 |f|_{\infty, X_G} .$$

By Theorem 1.2, there is a unique $\tilde{f} \in \mathcal{O}_{A\tau}(Y_G) \cap \mathcal{O}_{c\tau}(Y_G)$ such that $\tilde{f}|_{X_G} = f$ and

$$\max\{|\tilde{f}|_{A\tau, Y_G}, |\tilde{f}|_{c\tau, Y_G}\} \leq a_4 \max\{|f|_{A\tau, X_G}, |f|_{c\tau, X_G}\} \text{ for some } a_4 > 0.$$

Combining these inequalities with (2.3) and (2.1) yields

$$\max_K |\tilde{f}| \leq a_5 |f|_{\infty, X_G},$$

with some $a_5 > 0$ depending on X_G, Y_G only. For a fixed $g \in G$ consider $(g^*f)(z) := f(g(z))$. As above, there is a unique function $\tilde{f}_g \in \mathcal{O}_{A\tau}(Y_G) \cap \mathcal{O}_{c\tau}(Y_G)$, $\tilde{f}_g|_{X_G} = g^*f$, such that

$$\max_K |\tilde{f}_g| \leq a_5 |f|_{\infty, X_G},$$

But according to (2.2) and (1.1), function $(g^*\tilde{f})(z) := \tilde{f}(g(z))$ belongs to $\mathcal{O}_{A\tau}(Y_G)$ and $(g^*\tilde{f} - \tilde{f}_g)|_{X_G} \equiv 0$. Thus by Theorem 1.2 we have $\tilde{f}_g = g^*\tilde{f}$. Since K is the fundamental compact, the above inequality for each $\tilde{f}_g$ implies that

$$|\tilde{f}|_{\infty, Y_G} \leq a_5 |f|_{\infty, X_G}. \quad (2.4)$$

We will prove now that $a_5 = 1$ which gives the required statement. Indeed, the same arguments as above show that for any integer $n \geq 1$ the function $(\tilde{f})^n$ is the unique extension of f^n satisfying (2.4):

$$|(\tilde{f})^n|_{\infty, Y_G} \leq a_5 |f^n|_{\infty, X_G}.$$

Thus

$$|\tilde{f}|_{\infty, Y_G} \leq \lim_{n \rightarrow \infty} (a_5)^{1/n} |f|_{\infty, X_G} = |f|_{\infty, X_G}$$

The proof of the theorem is complete. $\square$

References

- [BD] D. E. Barrett and J. Diller, A new construction of Riemann surfaces with corona. J. Geom. Anal. **8** (1998), 341-347.
- [C] L. Carleson, Interpolation of bounded analytic functions and the corona problem. Ann. of Math. **76** (1962), 547-559.
- [G] T. W. Gamelin, Uniform algebras and Jensen measures. London Math. Soc. Lecture Notes Series **32**, Cambridge University Press, 1978.
- [GR] H. Grauert and R. Remmert, Theorie der Steinschen Räume. Springer-Verlag, Berlin, 1977.
- [JM] P. Jones and D. Marshall, Critical points of Green's functions, harmonic measure and the corona theorem. Ark. Mat. **23** no.2 (1985), 281-314.
- [L] F. Lárusson, Holomorphic functions of slow growth on nested covering spaces of compact manifolds. Canad. J. Math. **52** (5) (2000), 982-998.
- [N] T. Napier, Convexity properties of coverings of smooth projective varieties. Math. Ann. **286** (1990), 433-479.